# DISCUSSION PAPERS IN ECONOMICS

Working Paper No. 23-05


!
!
!
! "#$%&'(()%
*+,-$.),("%/0%1/#/.23/%&/'#3$.%
%
%
%

Current estimators that allow for selection based on a factor model either require (i) the number of time periods available is large, e.g. synthetic control (Abadie, 2021), factor-model imputation (Xu, 2017; Gobillon and Magnac, 2016), and the matrix completion method (Athey et al., 2021; Fernández-Val et al., 2021); or (ii) that an individual's error term, $u_{it}$, is uncorrelated over time (Imbens et al., 2021).[4] Both of these restrictions are non-realistic in many applied microeconomic data sets where the number of time periods is much smaller than the number of units and serial correlation of shocks is expected. Further, large-$T$ estimators often place restrictions on the dynamic heterogeneity of treatment. Our method requires neither large-$T$ or error term restrictions, but can still(but)-68ms,

our application and Section 6 leaves with some concluding remarks.

## 2 Model and Identification

We assume a panel data set with units $i = 1, \ldots, N$ and periods $t = 1, \ldots, T$. Treatment turns on in different periods for different units; we denote these groups by the period they start treatment. For each unit, we define $G_i$ to be unit $i$'s group with possible values $\{g_1, \ldots, g_G\}$ $G \subseteq \{2, \ldots, T\}$. We follow Callaway and Sant'Anna (2021) and denote $G_i = \infty$ for units that never receive treatment. We assume that $0 < P(G_i = g) < 1$ for all $g \in G \cup \{\infty\}$, so that the number of individuals in each group and the never-treated group grow with $N$. Treated potential outcomes are a function of group-timing, which we denote $y_{it}(g)$. For treatment indicators, we define the vector of treatment statuses $D_i = (D_{i1}, \ldots, D_{iT})$ where $D_{it} = 1(t \geq G_i)$ and the indicator $D_{ig} = 1(G_i = g)$ if unit $i$ is a member of group $g$. Let $T_0 = \min_j \{g_j\} - 1$ be the last period before the earliest treatment adoption.

Following Callaway and Sant'Anna (2021), we aim to estimate group-time average treatment effects on the treated:

$$ATT(g,t) = E(y_{it}(g) \mid G_i = g) - E(y_{it}(\infty) \mid G_i = g) \tag{2}$$

These quantities represent the average effect of treatment at time $t$ for units that start treatment in period $g$ for $t \geq g$. It is trivial to estimate other averages as well in our framework, including averaging over all post-treatment observations to estimate an overall $ATT$, and averaging over $(i,t)$ where $t - G_i = \ell$ to estimate event-study estimands $ATT(\ell)$'s. We discuss these and other extensions from Callaway and Sant'Anna (2021) in Section 3.

We now state our main identifying assumptions.

Assumption 1 (Sampling). The random vectors $\{(y_{it}, \ldots$